



CONSULTANTS IN ENGINEERING,
ENVIRONMENTAL SCIENCE &
PLANNING

BASELINE EMISSIONS INVENTORY REPORT

Baseline Emissions Inventory Report for County Offaly

Prepared for:
Offaly County Council




Date: January 2023

Core House, Pouladuff Road, Cork, T12 D773, Ireland
T: +353 21 496 4133 | E: info@ftco.ie

CORK | DUBLIN | CARLOW

www.fehilytimoney.ie



Baseline Emissions Inventory Report for County Offaly

REVISION CONTROL TABLE, CLIENT, KEYWORDS AND ABSTRACT

User is responsible for Checking the Revision Status of This Document

Rev. No.	Description of Changes	Prepared by:	Checked by:	Approved by:	Date:
00	Draft for Client Review	DH/RD/EW/BF/AMW/SJ/NSC	RD	BG	01/02/2023

Client: Offaly County Council

Keywords: Baseline emissions, greenhouse gas, carbon dioxide, methane, Offaly.

Abstract: This report presents the findings of a baseline emissions inventory study carried by Fehily Timoney and Company (FT) on behalf of Offaly County Council. The report presents a 'helicopter overview' of baseline greenhouse gas emissions from various sectors in a year. This baseline report aims to raise awareness of climate change and the impact that different sectors in the midlands region have on Ireland's overall carbon emissions. It provides Offaly County Council with the necessary information to make informed decisions on climate change actions to lower the county's carbon emissions. The sectors that have been included in this analysis are residential, commercial and industrial, industrial processes, agriculture, transport, waste and wastewater, and land use, and land use change and forestry (LULUCF). Emissions associated with the local authority's own operations have also been accounted for.

TABLE OF CONTENTS

1. INTRODUCTION	1
2. CONTEXT	2
2.1.1 Climate Change Challenges	2
2.1.2 2030 Energy and Emission Targets.....	2
2.1.3 Baseline Emissions Inventories - <i>Setting Local and Regional Targets</i>	3
3. BASELINE EMISSIONS INVENTORY.....	4
3.1 Residential	4
3.1.1 Methodology.....	4
3.1.2 Results	4
3.1.3 Assumptions and Limitations	6
3.1.4 Data Sources.....	6
3.2 Commercial and Industrial	7
3.2.1 Methodology.....	7
3.2.2 Results	7
3.2.3 Assumptions and Limitations	9
3.2.4 Data Sources.....	9
3.3 Industrial Processes.....	9
3.3.1 Methodology.....	9
3.3.2 Results	9
3.3.3 Assumptions and Limitations	10
3.3.4 Data Sources.....	11
3.4 Agriculture	11
3.4.1 Methodology.....	11
3.4.2 Results	12
3.4.3 Assumptions and Limitations	14
3.4.4 Data Sources.....	14
3.5 Transport	14
3.5.1 Methodology.....	14
3.5.2 Results	14
3.5.3 Assumptions and Limitations	16
3.5.4 Data Sources.....	16
3.6 Waste and Wastewater	17

3.6.1	Methodology	17
3.6.2	Results	17
3.6.3	Assumptions and Limitations	19
3.6.4	Data Sources.....	19
3.7	Land Use, Land Use Change and Forestry (LULUCF).....	19
3.7.1	Methodology.....	19
3.7.2	Results	20
3.7.3	Assumptions and Limitations	21
3.7.4	Data Sources.....	21
3.8	Local Authority	21
4. OVERVIEW AND MAIN CONCLUSIONS		23

LIST OF FIGURES

	<u>Page</u>
Figure 3-1: Overview of Residential Emissions.....	5
Figure 3-2: Breakdown of Main Space Heating Fuel for Dwellings	5
Figure 3-3: Percentage Breakdown of GHG emissions in the Commercial and Industrial Sector (combustion versus electricity)	8
Figure 3-4: Percentage Breakdown of combustion GHG emissions (commercial services versus manufacturing combustion).....	8
Figure 3-5: Breakdown of Industrial Processes Emissions	10
Figure 3-6: Percentage Breakdown of Livestock Emissions	13
Figure 3-7: Percentage Breakdown of Agricultural Emissions Overall	13
Figure 3-8: Breakdown of Transport Emissions in the County	15
Figure 3-9: Emissions per Single Unit of Vehicle Type.....	16
Figure 3-10: Emission from the Waste Sector in the County	18
Figure 3-11: Breakdown of LULUCF Emissions in the County	20
Figure 3-12: Percentage Breakdown of GHG Emissions for the Local Authority	22
Figure 4-1: Breakdown of Emissions Per Sector in the County	24
Figure 4-2: Emission Reduction Requirements Per Sector.....	25

LIST OF TABLES

	<u>Page</u>
Table 3-1: Overview of Residential Emissions.....	4
Table 3-2: Breakdown of Industrial Processes Emissions	10
Table 3-3: Breakdown of Livestock Numbers in the County	12
Table 3-4: Breakdown of Agricultural Emissions.....	12
Table 3-5: Breakdown of Transport Emissions in the County	15
Table 3-6: Emissions from the Waste Sector in the County.....	18
Table 3-7: Wastewater Emissions	19
Table 3-8: Breakdown of LULUCF Emissions in the County	20
Table 3-9: GHG Emissions Data for the Local Authority.....	21
Table 4-1: Breakdown of Emissions Per Sector in the County	23



1. INTRODUCTION

This report presents the findings of a 'Tier 1' Baseline Emission Inventory (BEI) study carried out by Fehily Timoney and Company (FT) for the county of Offaly. The study has been carried out on behalf of Offaly County Council. The report evaluates and determines the baseline greenhouse gas (GHG) emissions from various societal sectors in the county in 2018 and allows Offaly County Council to measure the emission reductions required to achieve the emission reduction target of reducing GHG emissions by 51% by 2030. This baseline report aims to raise awareness of climate change and the impact that different sectors in the midlands region have on Ireland's overall carbon emissions. It provides Offaly County Council with the necessary information to make informed decisions on climate change actions to lower the county's carbon emissions.

The sectors that have been included in this analysis residential, commercial and industrial, industrial processes, agriculture, transport, waste and wastewater, and land use, and land use change and forestry (LULUCF).

The methodology is based on the Tier 2 'Bottom Up' Approach defined in Annex C to the Draft Local Authority Climate Action Plan Guidelines.

The national emission reduction target of 51% by the end of 2030 is based on the GHG emissions reported for the end of 2018, in the national GHG emissions inventory. Accordingly, the data collated and analysed to inform this BEI is relative to the baseline year of 2018, or the nearest year possible to 2018.

GHG emissions are reported as Carbon dioxide (CO₂) or Carbon dioxide equivalent (CO₂-eq) in this report. The Global Warming Potential (GWP) of other GHGs (e.g., Methane, CH₄) has been factored to allow for reporting in CO₂-eq. Where the term 'emissions' in and of itself is used in this report, this refers solely to GHG emissions, and not non-GHG emissions (such as Sulphur oxides, SO_x, or Nitrogen oxides, NO_x).



2. CONTEXT

2.1.1 Climate Change Challenges

Climate change refers to the long-term changes in the earth's weather patterns or average temperatures. In Ireland this is demonstrated by rising sea levels, extreme weather events and changes in the eco-system. Extensive research and a significant body of evidence has shown a correlation between the increasing global average temperature and the increasing quantity of GHG released into the atmosphere, particularly from anthropogenic sources.

Changes in weather patterns and climate can have significant adverse impacts on the environment and human beings. The Intergovernmental Panel on Climate Change (IPCC) published the *Climate Change 2022: Impacts, Adaptation and Vulnerability in 2022*. Included in this report is an outline of observed impacts of climate change on the environment and human beings. These include impacts from inland flooding, damages to infrastructure, impacts from infectious disease, displacement, animal and livestock health and productivity, mental health and water scarcity derived from climate change.

The seriousness of the potential impacts and risks associated with climate change is reflected in the vast quantity of legislation that has been introduced to mitigate those impacts and risks, beginning with the establishment of the United Nations Framework Convention on Climate Change (UNFCCC) treaty in 1992. Subsequent significant, international treaties, agreements have been made and international gathering taken place since, including, the Kyoto Protocol, the Paris Agreement, 2030 Climate and Energy Policy Framework and the twenty-seven UNFCCC, Convention of the Parties (COP) conferences that have taken places since its formation.

The Paris Agreement was the first-ever universal, legally binding global climate change agreement, adopted at the Paris climate conference (COP21) in December 2015.

There are many significant additional benefits to reducing GHG emission levels and increasing the share of renewable energies. These include a decrease in dependency on fossil fuels, which in turn results in a higher security of energy supply, better health, lower energy costs, an increase in the county's competitiveness, and a more sustainable economy.

2.1.2 2030 Energy and Emission Targets

The EPA has confirmed that Ireland has exceeded its 2020 target for GHG emissions by 6.73 Mt CO₂-eq. Projections indicate that Ireland can meet its targets over the 2021 – 2030 period but only with the full implementation of the Climate Action Plan and use of flexibilities the European Effort Sharing Regulation (ESR) provides.

The Climate Action and Low Carbon Development Act 2015 specifies that a national low carbon transition and mitigation plan and sectoral adaptation plans be drafted and approved by the Government in relation to climate change for the purpose of pursuing the transition to a low carbon, climate resilient and environmentally sustainable economy. The act required the establishment of the Climate Change Advisory Council and the creation and approval by the government of a National Mitigation Plan (to be published every five years), National Adaptation Framework and an Annual Transition Statement.



In 2017, the first National Mitigation Plan for Ireland was published but replaced in 2019 with the publication of a Climate Action Plan (Government of Ireland, 2019). In 2021, this Climate Action Plan (CAP) was subsequently updated. The 2021 plan sets out actions that must be taken to ensure delivery of commitments in the Programme of Government and the Climate Act 2021 (e.g., reducing GHG emissions by 51% by 2030 and to achieve net zero emissions by 2050). The achievement of this target will require significant, unanimous effort across all government departments and local authorities.

2.1.3 Baseline Emissions Inventories - Setting Local and Regional Targets

A key element of the Climate Action and Low Carbon Development Act as amended, relevant to local authorities is the requirement for local authorities to prepare individual climate action plans (CAPs) for their functional area. The plans include climate mitigation and adaptation measures and are required to be updated every five years. Local Authority Development Plans must also be aligned with their CAP.

The primary goal of the national CAP is to ensure a just transition to a climate neutral Ireland using ‘*an integrated, structured and evidence-based approach.*’ The CAP also states, ‘*our capacity to anticipate and plan for a just transition requires the development of a robust evidence base to support policy development and effective ongoing monitoring.*’ Furthermore ‘*data will play a critical role in assisting local authorities in forecasting where changes will occur (or are already underway), and who will be most impacted.*’

The primary objective of this report is the generation of BEI for the local authority functional area of County Offaly. The BEIs will be a contemporary snapshot in time of emissions in County Offaly and will act as a reference point on which specific county wide and/or sectoral targets can be established to manage and reduce GHG emissions.

The BEIs will also allow the monitoring and tracking of progress towards meeting those targets and the efficacy of any measures introduced by the local authority, through future updating of the emissions inventories.

Assessing sectoral contributions will define where reductions in GHG emissions are most needed to achieve GHG emissions targets. By assessing sectoral contributions to climate change and the risks they pose with respect to climate change, local authorities can rank sectors and sub-sectors with respect to their counties contribution to climate change and make informed, practical decisions on how they can contribute to and influence mitigating these risks within each sector in partnership with relevant stakeholders.

CAPs will be established for a duration of five years. It is expected that, at a minimum, and to maintain an up-to-date and accurate evidence base to inform decision making, BEIs would be revised at least every five years in line with each revision of the local authority CAPs. However, it is also advised that BEIs should be updated when new, significant evidence, data etc. is available.



3. BASELINE EMISSIONS INVENTORY

3.1 Residential

3.1.1 Methodology

The following methodology was used to determine emissions from the Residential sector.

- Map Elre on combustion related GHG emissions for the residential sector in the county was sourced.
- Central Statistics Office (CSO) data on metered electricity consumption for the residential sector in the county was sourced. GHG emissions associated with electricity consumption were calculated using the Sustainable Energy Authority Ireland (SEAI) emission factor for electricity in 2018.
- Residential sector GHG emissions for the county and per capita were calculated having regard to the above data.
- An estimate for main source of heating fuel for dwellings in the county was made using data from the CSO, which in turn is sourced from the Building Energy Rating (BER) database. This data was used to determine the number of dwellings in the county that rely on the following fuel types as their main source of space heating: Mains Gas, LPG, Heating Oil, Electricity, Solid Fuel. This aids the characterization of Residential sector emissions, providing an additional layer of insight.

3.1.2 Results

A breakdown of residential sector emission for the county for the baseline year is presented in Table 3-1 and Figure 3-1.

Table 3-1: Overview of Residential Emissions

Combustion Emissions in the County (from space heating and hot water) tCO ₂ -eq	187,786.15	74.52%
Electricity Emissions in the County (from electrical appliances and lighting) tCO ₂ -eq	64,212	25.48%




Figure 3-1: Overview of Residential Emissions

The majority of residential sector emissions are from the combustion of fuels for space heating and hot water (75%). Electricity related emissions account for 26% of residential emissions in the county.

The calculated emissions per capita in the baseline year for the county was 3.23 tCO₂-eq. This is a high figure relative to other counties in the Midlands region.

For an additional level of insight, a breakdown of main space heating fuels used at dwellings in the county is provided in Figure 3-2.




Figure 3-2: Breakdown of Main Space Heating Fuel for Dwellings



Heating oil is the primary source of main space heating fuel in the county (56%). It is commonly used in detached and semi-detached dwellings in rural areas. The combustion of heating oil generates a higher level of emissions per unit of energy compared to the combustion of many other heating fuels such as gas or LPG. The use of heating oil has significant contribution to residential sector emissions in the county.

Electricity accounts for 15% of main space heating fuel use in the county. This is a relatively high figure. This practice is generally more common in apartments.

Mains gas accounts for 14% of main space heating fuel use in the county. This is due to presence of a gas network in parts of the county.

A significant number of dwellings use solid fuel as a main space heating fuel (14%). This is likely due to the presence of a significant level of bogland in the county and the relatively high number of dwellings that use peat sourced from these lands as their main space heating fuel. Solid fuel burning will generate a relatively higher level of emissions compared to other commonly used space heating fuels due to its carbon intensity.

The high level of combustion related emissions in the county is likely to be accounted for in part by the practice of solid fuel burning.

3.1.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the Residential sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.
- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.
- The CSO database for heating fuel type only considers dwellings in the county that have had BER Ratings completed from them. The use of BER data is limited in that it only provides data for dwellings that have had BERs carried out on them, which only represents a proportion of the general housing stock. BER data does however provide a good representation of the housing stock generally given the level of BERs carried out on housing in the county.

3.1.4 Data Sources

1. Map Elre / Environmental Protection Agency National Emission Inventory, Local Authority Emissions, 2021.
2. Central Statistics Office, Metered Electricity Consumption, 2018.
3. Central Statistics Office, Main Space Heating Fuel Per County, 2018.
4. SEAI, Energy in Ireland, 2018.
5. CSO, Census of Population, 2016.



3.2 Commercial and Industrial

3.2.1 Methodology

The following methodology was used to determine emissions from the Commercial and Industrial sector.

- Raw data on commercial and industrial sector combustion related GHG emissions for 2018 was sourced from the Map Elre / the EPA's National Emission Inventory (2021) for the county. This emissions data is broken down by emissions from the combustion of fossil fuels in the commercial services sub-sector and the manufacturing sub-sector.
- Total non-residential electricity use levels for 2018 for the county was sourced from Electricity Supply Board (ESB) / CSO databases on metered electricity consumption. This total is then multiplied by the SEAI's electricity emission factor for 2018 to determine electricity related GHG emissions in the commercial and industrial sector for the year.
- The total GHG emissions in tCO₂ for the county is then calculated by adding the combustion and electricity GHG emissions.

3.2.2 Results

The commercial and industrial sector in the county generates 100,043 tCO₂. 65% of these emissions are due to electricity consumption in the sector, whilst 35% are due to combustion related emissions (See Figure 3-3).

53% of combustion related emissions originate from the manufacturing sub-sector. 47% of combustion related emissions originate from the commercial services sub-sector. (See Figure 3-4).

This sector in the county has a higher proportion of combustion related emissions in comparison to most other counties in the midlands. A higher proportion of these combustion emissions originate from manufacturing.




Figure 3-3: Percentage Breakdown of GHG emissions in the Commercial and Industrial Sector (combustion versus electricity)




Figure 3-4: Percentage Breakdown of combustion GHG emissions (commercial services versus manufacturing combustion)



3.2.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the Commercial and Industrial sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.
- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.

No other assumptions and limitations exist in relation to the Commercial and Industrial sector data. It is noted that 'Industrial Processes' is considered as a separate category within this report.

3.2.4 Data Sources

- Map Elre / Environmental Protection Agency, National Emission Inventory, Local Authority Emissions, 2021.
- MEC03 - Metered Electricity Consumption. Central Statistics Office. <<https://data.cso.ie/>>. Accessed on the 23rd November 2022.
- Energy in Ireland – 2019 Report. Sustainable Energy Authority of Ireland, 2019. <<https://www.seai.ie/publications/Energy-in-Ireland-2019-.pdf>>

3.3 Industrial Processes

3.3.1 Methodology

The following methodology was used to determine emissions from the Industrial Processes sector.

- A breakdown of emissions for the sector was obtained from the Map Elre database. The Access database file contains emissions from all sectors for all Irish counties separated into different sub-categories. The emissions data for Industrial Processes specific to the county was extracted.
- The GHG Emissions from the database are categorised into various pollutants (i.e. CO₂, CH₄, N₂O and SF₆). The emissions figures for these pollutants were converted to CO₂-eq with reference to the GWP value for each substance.

3.3.2 Results

A breakdown of industrial process related emissions is provided in Table 3-2 and illustrated in Figure 3-5.



Table 3-2: Breakdown of Industrial Processes Emissions

Industry Category	GHG Emissions (tCO ₂ eq)
Mineral	238.37
Chemical	0.00
Metal	0.00
Non-energy Products from Fuels and Solvent Use	1,876.12
Electronics	0.00
Product Uses as Substitutes for ODS	0.00
Other Product Manufacture and Use	7,552.32
Other - Food and Beverage	120.60




Figure 3-5: Breakdown of Industrial Processes Emissions

GHG emissions from industrial processes in the county arise from 4 main subcategories. Emissions from 'Other Product Manufacture and Use' are the highest (77%) at 7,552 tonne CO₂eq, followed by 'Non-energy products from fuels and solvent use' (19%) at 1,876 tonne CO₂eq. A relatively minor level of emissions arise from 'Minerals' (3%) and 'Other - Food and Beverage' (1%).

3.3.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the Industrial Processes sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high-level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.



- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.

3.3.4 Data Sources

- Map Elre / Environmental Protection Agency National Emission Inventory 2021.

3.4 **Agriculture**

3.4.1 Methodology

County level agricultural data was obtained from the CSO's Census of Agriculture. This data was broken down into different agricultural activities relevant to the county.

Once all data was obtained, a breakdown of agricultural sector related emissions was developed. These emissions were broken into two categories, as follows:

- Livestock (sum of cattle (suckler and dairy), pigs, sheep and poultry).
- Managed Soils (sum of direct and indirect N₂O emissions, limestone emissions and urea application emissions).

Livestock emissions at county level was determined with reference to national emissions statistics, the national herd, and the herd in the county. This was considered to be the most representative and accurate method for determining emissions for livestock.

Managed soils emissions at county level were estimated with reference to national emission statistics for direct and indirect N₂O emissions, limestone emissions and urea application emissions; and the area of managed agricultural soils nationally compared to the area of managed soils in the county.



3.4.2 Results

A breakdown of livestock numbers in the county is presented in Table 3-3.

Table 3-3: Breakdown of Livestock Numbers in the County

Breakdown of Livestock Numbers in the County	
Livestock Type	Livestock Numbers
Dairy Cows	35,738
Other Cattle and Cows	218,950
Sheep	103,701
Pigs	109,872
Poultry	55,083

Cattle farming is the primary type of livestock farming in the county. Other cattle and cows (for beef production) constitute 42% of total livestock numbers within the county. In terms of numbers, this is followed by pigs, then sheep, poultry and dairy cows.

A breakdown of agricultural related emissions in the baseline year is presented in Table 3-4.

Table 3-4: Breakdown of Agricultural Emissions

Type	CO ₂ -eq Emissions (tCO ₂ -eq)
Livestock	
Cattle	459,602.15
Sheep	14,979.72
Pigs	22,938.77
Poultry	522.05
Managed Soils	
Managed Soils	188,635
Totals	
Total Livestock	498,042.70
Total Managed Soils	188,634.60
Overall Total	686,677.30

A breakdown of livestock related emissions is presented in Figure 3-6.




Figure 3-6: Percentage Breakdown of Livestock Emissions

Emissions from cattle (beef and dairy cows) combined contribute the most in terms of livestock related emissions by far, accounting for 92% of CO₂-eq emissions from agriculture in the county. Cattle produce much more methane emissions than other livestock, and due to the higher global warming potential of methane, their impact is much larger on the environment. Emissions associated with sheep, pig and poultry farming are relatively low by comparison.

Managed soils in the county generated 188,634.60 tCO₂-eq emissions in the baseline year.

Agricultural emissions for the county broken down by the two overarching categories, total livestock, and managed soils combined are presented in Figure 3-7.




Figure 3-7: Percentage Breakdown of Agricultural Emissions Overall



Livestock emissions account for the majority of emissions from the sector overall (73%). Managed soils emissions are lower by comparison, yet still significant (27%).

3.4.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the Agricultural sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.
- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.
- National emissions data for 'Other Livestock' defined in the EPA's national emission inventory has been assumed to relate to Poultry. This 'Other Livestock' category includes poultry, goats, horses and mink, however, in terms of population, poultry accounts for the vast majority of livestock under this category.
- It is assumed that all managed agricultural soils in the county release the same level of emissions. In reality, emissions levels will vary depending on the type and level of fertilizer, lime or urea application on those soils based on their specific use (i.e., pasture, wheat, barley, potatoes).

3.4.4 Data Sources

- Map Elre / Environmental Protection Agency, National Emission Inventory 2021.
- Central Statistics Office, Census of Agriculture, 2020.

3.5 Transport

3.5.1 Methodology

The following methodology was applied to determine transport related emissions in the county:

- GHG emissions data for the transport sector was sourced from the Map Elre database / the EPA's National Emission Inventory 2021. This database provides a breakdown of emission for a number of transport sub-categories. Various emission types (CH₄, N₂O) reported were converted to tCO₂-eq using the GWP for each type of emission.
- For an additional layer of insight, an estimation of GHG emissions per single unit of each vehicle type is made by dividing emissions associated with each vehicle type by vehicle population. The CSO's Transport Omnibus from 2018 was reviewed to ascertain the vehicle population in the county for 2018 for each vehicle type.

3.5.2 Results

Detail on GHG emissions associated with each transport category in the county is provided in Table 3-5 and illustrated in Figure 3-8.



Table 3-5: Breakdown of Transport Emissions in the County

Transport Category	GHG Emissions (tCO ₂ -eq)
Passenger cars	136,033.34
Light-duty trucks	36,479.02
Heavy-duty vehicles	73,268.62
Motorcycles	315.31
Railways	5,006.06




Figure 3-8: Breakdown of Transport Emissions in the County

Passenger cars are the primary source of transport related GHG emission in the county (54%), followed by heavy duty vehicles (29%), then light duty trucks (15%). Emissions from railways are relatively minor, which reflects the low level of railway services in the county. Emissions from motorcycles are relatively minor, which reflects the low number of this type of vehicle in the county. Transport related emission in the county were closely similar to other counties in the Midlands region

An estimation of emissions per single unit of each vehicle type is presented in Figure 3-9.




Figure 3-9: Emissions per Single Unit of Vehicle Type

Goods vehicles are the most carbon intensive vehicle type in the county. This reflects the relatively high level of carbon emissions associated with these weight laden vehicles. Private cars are next most carbon intensive vehicle type in the county, followed by motorcycles.

3.5.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the Transport sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.
- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.
- When determining emissions per single unit of vehicle type goods vehicles (as defined by the CSO) were assumed to be inclusive of both light duty trucks and heavy duty vehicles. Heavy duty vehicles were also inclusive of buses.

3.5.4 Data Sources

- Map Elre / Environmental Protection Agency, National Emission Inventory 2021.
- Transport Omnibus 2018, Central Statistics Office, 2018.



3.6 Waste and Wastewater

3.6.1 Methodology

3.6.1.1 *Waste*

Qualitative analysis of the waste sector in the county was carried out to determine the level of GHG emissions associated with the sector. The following waste categories defined in the EPA's National Emission Inventory 2021 were examined.

- Managed Waste Disposal,
- Composting,
- Anaerobic Digestion (AD),
- Incineration, and
- Open Burning of Waste.

Using national emissions data for the managed waste disposal, incineration and open burning of waste categories defined in the EPA's National Emission Inventory 2021, emissions associated with this sector for the county were estimated for the county on a pro-rata basis considering national and county population levels. The accumulated emissions data for these waste activities in the county were combined to determine emissions in tCO₂-eq for the baseline year.

There are no composting or anaerobic digestion facilities in the county. Thus, there are no emission in the county from these categories of waste facility.

For open burning of waste, there is no local data on GHG emissions from this unregularized and uncontrolled activity.

3.6.1.2 *Wastewater*

Using national emissions data for the wastewater sector defined in the EPA's National Emission Inventory 2021, emissions associated with this sector for the county were estimated for the county on a population pro-rata basis.

3.6.2 Results

3.6.2.1 *Waste*

A breakdown of the emissions from the waste sector in the county is presented in Table 3-6 and illustrated in Figure 3-10.



Table 3-6: Emissions from the Waste Sector in the County

Waste Category	National GHG Emissions 2018 (tCO ₂ eq)	County GHG Emissions (tCO ₂ eq)
Managed Waste Disposal Sites	692,700	11,341
Waste incineration	20,300	332
Open burning of waste	3,630	59
Composting	42,000	0
Anaerobic Digestion at biogas facilities	2,400	0




Figure 3-10: Emission from the Waste Sector in the County

The waste sector in the county generated 11,733 tCO₂ in the baseline year. 97% of waste sector emissions originate from managed waste disposal sites, namely managed landfills including closed or historic landfills still generating Methane and Carbon dioxide emissions due to the decomposition of waste inside the waste bodies of the landfills.

3.6.2.2 Wastewater

Estimates for GHG emissions from the wastewater sector in the county for the baseline year are presented in Table 3-7.



Table 3-7: Wastewater Emissions

Sector Category	National GHG Emissions 2018 (tCO ₂ eq)	County GHG Emissions 2018 (tCO ₂ eq)
Domestic Wastewater	147,900.00	2,421.4

3.6.2.3 Summary

A summary of the key findings for the waste and wastewater sectors is presented below:

- Total emissions from the waste sector in the county were estimated to be 11,733 tCO₂-eq.
- The majority of waste sector emissions in the county are generated by managed landfills.
- Total emissions from the wastewater sector in the County were estimated to be 2,421.4 tCO₂-eq, which represents a small fraction of domestic wastewater emissions nationally.

3.6.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the Waste and Wastewater sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high-level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.
- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.
- Managed waste disposal related to all regularized landfills including operational landfills and closed or historic landfills.

3.6.4 Data Sources

- Map Elre / Environmental Protection Agency, National Emission Inventory, EPA, 2021.
- EPA IE licence, IPC licence, waste licence and WFP search databases and licence files for landfills contained within them - accessed 28/11/2022.

3.7 Land Use, Land Use Change and Forestry (LULUCF)

3.7.1 Methodology

The following methodology was used to determine emissions from the LULUCF sector.

- A breakdown of emissions for the sector was obtained from the Map Elre database. The Access database file contains emissions from all sectors for all Irish counties separated into different sub-categories. The emissions data for LULUCF specific to the county was extracted.



- The GHG Emissions from the database are categorised into various pollutants (i.e. CO₂, CH₄, N₂O and SF₆). The emissions figures for these pollutants were converted to CO₂-eq with reference to the GWP value for each substance.

3.7.2 Results

A breakdown of GHG emissions relating to land use, land use change and forestry for the various land uses in the county for the baseline year is presented in Table 3-8 and Figure 3-11.

Table 3-8: Breakdown of LULUCF Emissions in the County

Sector Category	GHG Emissions (tCO ₂ eq)
Forestland	-169,925.04
Cropland	-2,370.41
Grassland	240,290.77
Wetlands	75,262.38
Settlements	4,945.61
Other land	0.00
Harvested wood products	-11,088.55




Figure 3-11: Breakdown of LULUCF Emissions in the County



Grassland in the county is the land use type that contributes most in terms of emissions, followed by the Wetlands, Settlements and Other Land categories. Forestland and Cropland serve to absorb CO₂ considering both CO₂ gains and losses overall. Forestland and Cropland generally absorb more carbon (e.g., through vegetative photosynthesis) than they release (e.g., through deforestation or harvesting). In terms of harvested wood products (HWP), after harvest, atmospheric carbon (C) is immediately sequestered by vegetation re-growth. This temporal mismatch between oxidation of HWPs and C uptake by vegetation generates a net sink.

3.7.3 Assumptions and Limitations

The following assumptions and limitations exist in relation to the LULUCF sector data:

- The 'Top Down' methodology employed in this instance results in a broad, high level estimation of GHG emissions from this sector in the county. A degree of 'uncertainty of measurement' therefore exists in relation to the emissions data.
- All assumptions and limitations relating to Map Elre national inventory emission data apply to the emissions data presented, as this data is informed by Map Elre data.

3.7.4 Data Sources

- Map Elre / Environmental Protection Agency, National Emission Inventory 2021.

3.8 Local Authority

As detailed in Annex C to the Draft Local Authority Climate Action Guidelines, public sector including local authority GHG emissions are captured under the sectors dealt with so far in this report (e.g., Institutional emissions are captured under the Commercial sector, Public Transport emissions are captured under the Transport Sector). The guidance advises that a separate BEI for local authority GHG emissions is prepared, however, having regard to the local authority's responsibility establish a GHG emission baseline and deliver on its own targets for GHG emission reductions.

GHG emissions data for the local authority was sourced from the county council's SEAI monitoring and reporting database. This data is presented in Table 3-9 and Figure 3-12.

Table 3-9: GHG Emissions Data for the Local Authority

Category	GHG Emission tCO ₂ -eq
Electricity	2,206
Thermal	698
Transport	643
Total	3,547




Figure 3-12: Percentage Breakdown of GHG Emissions for the Local Authority



4. OVERVIEW AND MAIN CONCLUSIONS

The analysed sectors in the county generated 1,450,877.1 tCO₂-eq in the baseline year, overall. A breakdown of these emissions and targeted emission levels for 2030 by sector is presented in Table 4-1.

Table 4-1: Breakdown of Emissions Per Sector in the County

Sector	2018 Emissions tCO ₂ -eq	Percentage Breakdown	Sectoral Emission Ceiling Reduction Percentage 2030	Target Emissions 2030 tCO ₂ -eq
Residential	251,999	17%	40%	151,199
Commercial and Industrial*	100,043	7%	45%	55,023
Industrial Processes	9,787	1%	35%	6,362
Agriculture	686,677	47%	25%	515,008
Transport	251,102	17%	50%	125,551
Waste and Wastewater	14,154	1%	50%	7,077
LULUCF **	137,115	9%	-	-
Total	1,450,877	100%	-	-

*Commercial and Industrial GHG emissions are unable to be disaggregated considering the methodology applied to calculate GHG emission for these sectors and the available data that underpins the calculations (e.g., electricity use data sourced from the CSO is only available for the 'non-residential sector' and does not differentiate between the commercial and industrial sector). As such, the more conservative 'Commercial' sectoral emission ceiling reduction has been applied.

** Finalization of the sectoral emission ceiling for the LULUCF sector has been deferred for up to 18 months from July 2022 to allow for the completion of a national land use review.

A percentage breakdown of emissions in the baseline year is provided in Figure 4-1.




Figure 4-1: Breakdown of Emissions Per Sector in the County

The top three sectors in the county in terms of GHG emission levels were Agriculture, Residential, and Transport producing 47%, 17% and 17% of tCO₂-eq respectively, of the total emissions in the county. From this analysis, these sectors should be the main targets of energy and emission initiatives.

An illustration of the target emissions for 2030 broken down by each sector is provided in Figure 4-2.




Figure 4-2: Emission Reduction Requirements Per Sector



**CONSULTANTS IN ENGINEERING,
ENVIRONMENTAL SCIENCE
& PLANNING**

www.fehilytimoney.ie

📍 Cork Office

Core House
Pouladuff Road,
Cork, T12 D773,
Ireland
+353 21 496 4133

📍 Dublin Office

J5 Plaza,
North Park Business Park,
North Road, Dublin 11, D11 PXT0,
Ireland
+353 1 658 3500

📍 Carlow Office

Unit 6, Bagenalstown Industrial
Park, Royal Oak Road,
Muine Bheag,
Co. Carlow, R21 XW81,
Ireland
+353 59 972 3800

